

Spin-asymmetry energy of nuclear matter

N. Kaiser

Physik-Department T39, Technische Universität München, D-85747 Garching, Germany

Abstract

We calculate the density-dependent spin-asymmetry energy $S(k_f)$ of isospin-symmetric nuclear matter in the three-loop approximation of chiral perturbation theory. The interaction contributions to $S(k_f)$ originate from one-pion exchange, iterated one-pion exchange, and (irreducible) two-pion exchange with no, single, and double virtual Δ -isobar excitation. We find that the truncation to 1π -exchange and iterated 1π -exchange terms (which leads already to a good nuclear matter equation of state) is spin-unstable, since $S(k_{f0}) < 0$. The inclusion of the chiral $\pi N\Delta$ -dynamics guarantees the spin-stability of nuclear matter. The corresponding spin-asymmetry energy $S(k_f)$ stays positive within a wide range of an undetermined short-range parameter S_5 (which we also estimate from realistic NN-potentials). Our results reemphasize the important role played by two-pion exchange with virtual Δ -isobar excitation for the nuclear matter many-body problem. Its explicit inclusion is essential in order to obtain good bulk and single-particle properties.

PACS: 12.38.Bx, 21.30.-x, 21.65.+f

In recent years a novel approach to the nuclear matter problem based on effective field theory (in particular chiral perturbation theory) has emerged. The key element there is a separation of long- and short-distance dynamics and an ordering scheme in powers of small momenta. At nuclear matter saturation density $\rho_0 \simeq 0.16 \text{ fm}^{-3}$ the Fermi momentum k_{f0} and the pion mass m_π are comparable scales ($k_{f0} \simeq 2m_\pi$), and therefore pions must be included as explicit degrees of freedom in the description of the nuclear many-body dynamics. The contributions to the energy per particle $\bar{E}(k_f)$ of isospin-symmetric (spin-saturated) nuclear matter as they originate from chiral pion-nucleon dynamics have been computed up to three-loop order in refs.[1, 2]. Both calculations are able to reproduce correctly the empirical saturation point of nuclear matter by adjusting one single parameter (either a coupling $g_0 + g_1 \simeq 3.23$ [1] or a cut-off $\Lambda \simeq 0.65 \text{ GeV}$ [2]) related to unresolved short-distance dynamics.¹ The novel mechanism for saturation in these approaches is a repulsive contribution to the energy per particle $\bar{E}(k_f)$ generated by Pauli-blocking in second order (iterated) one-pion exchange. As outlined in section 2.5 of ref.[2] this mechanism becomes particularly transparent by taking the chiral limit $m_\pi = 0$. In that case the interaction contributions to $\bar{E}(k_f)$ are completely summarized by an attractive k_f^3 -term and a repulsive k_f^4 -term where the parameter-free prediction for the coefficient of the latter is very close to the one extracted from a realistic nuclear matter equation of state.

In a recent work [3] we have extended the chiral approach to nuclear matter by including systematically the effects from two-pion exchange with single and double virtual $\Delta(1232)$ -isobar excitation. The physical motivation for such an extension is threefold. First, the spin-isospin- $3/2$ $\Delta(1232)$ -resonance is the most prominent feature of low-energy πN -scattering. Secondly, it

¹The cut-off scale Λ serves the purpose to tune the strength of an attractive zero-range NN-contact interaction.

is well known that the two-pion exchange between nucleons with excitation of virtual Δ -isobars generates the needed isoscalar central NN-attraction [4] which in phenomenological one-boson exchange models is often simulated by a fictitious scalar " σ "-meson exchange. Thirdly, the delta-nucleon mass splitting of $\Delta = 293$ MeV is of the same size as the Fermi momentum $k_{f0} \simeq 2m_\pi$ at nuclear matter saturation density and therefore pions and Δ -isobars should both be treated as explicit degrees of freedom. A large variety of nuclear matter properties has been investigated in this extended framework in ref.[3]. It has been found that the inclusion of the chiral $\pi N\Delta$ -dynamics is able to remove most of the shortcomings of previous chiral calculations of nuclear matter [2, 5, 6, 7]. For example, the momentum-dependence of the (real) single-particle potential $U(p, k_f)$ near the Fermi surface $p = k_f$ improves significantly. As a consequence of that the critical temperature of the first-order liquid-gas phase transition of isospin-symmetric nuclear matter gets lowered to the realistic value $T_c \simeq 15$ MeV. The isospin properties of nuclear matter improve also substantially when including the chiral $\pi N\Delta$ -dynamics. Instead of bending downward above ρ_0 as in previous chiral calculations [2, 7], the energy per particle of pure neutron matter $\bar{E}_n(k_n)$ and the (isospin) asymmetry energy $A(k_f)$ grow now monotonically with density (see Figs. 10,11 in ref.[3]). Good agreement with results of sophisticated many-body calculations and (semi)-empirical values has been found in the density regime $\rho = 2\rho_n < 0.2 \text{ fm}^{-3}$ relevant for conventional nuclear physics.

Given all that success of chiral perturbation theory for the nuclear matter many-body problem it is nevertheless still necessary to check the spin-stability of nuclear matter in that framework. Such an investigation is the subject of the present paper. We remind that in the past the criterion of spin-stability of infinite nuclear matter has required modifications of several phenomenologically very successful Skyrme forces [8, 9, 10]. For recent work on generalized symmetry energy coefficients in the context of phenomenological Skyrme forces, see also ref.[11].

Let us begin with defining the spin-asymmetry energy $S(k_f)$ of infinite nuclear matter. Consider this homogeneous many-nucleon system in a partially spin-polarized state such that the Fermi seas of the spin-up and spin-down nucleons are filled unequally high. With the help of the spin-projection operators $(1 \pm \sigma_3)/2$ such a spin-unsaturated configuration is realized by the substitution:

$$\theta(k_f - |\vec{p}|) \rightarrow \frac{1 + \sigma_3}{2} \theta(k_\uparrow - |\vec{p}|) + \frac{1 - \sigma_3}{2} \theta(k_\downarrow - |\vec{p}|), \quad (1)$$

in the medium insertion.² Here, $k_\uparrow$ and $k_\downarrow$ denote the different Fermi momenta of the spin-up and spin-down nucleons. Choosing $k_{\uparrow,\downarrow} = k_f(1 \pm \eta)^{1/3}$ (with η a small parameter) the total nucleon density $\rho = (k_\uparrow^3 + k_\downarrow^3)/3\pi^2 = 2k_f^3/3\pi^2$ stays constant. The expansion of the energy per particle of spin-polarized (isospin-symmetric) nuclear matter:

$$\bar{E}_{\text{pol}}(k_\uparrow, k_\downarrow) = \bar{E}(k_f) + \eta^2 S(k_f) + \mathcal{O}(\eta^4), \quad k_{\uparrow,\downarrow} = k_f(1 \pm \eta)^{1/3}, \quad (2)$$

around the spin-saturation line ($k_\uparrow = k_\downarrow$ or $\eta = 0$) defines the spin-asymmetry energy $S(k_f)$. The obvious criterion for the spin-stability of nuclear matter is then the positivity of the spin-asymmetry energy, $S(k_f) > 0$. The energy per particle at fixed nucleon density ρ must take on its absolute minimum value in the spin-saturated configuration.

The first contribution to the spin-asymmetry energy $S(k_f)$ comes from the kinetic energy $\sqrt{M^2 + p^2} - M$ of a non-interacting relativistic Fermi gas:

$$S(k_f) = \frac{k_f^2}{6M} - \frac{k_f^4}{12M^3}, \quad (3)$$

²Medium insertion is a technical notation for the difference between the in-medium and vacuum nucleon propagator. For further details, see section 2 in ref.[2].




Figure 1: The two-loop one-pion exchange Fock diagram and the three-loop iterated one-pion exchange Hartree and Fock diagrams. The combinatoric factors of these diagrams are $1/2$, $1/4$ and $1/4$, in the order shown. Their isospin factors for isospin-symmetric nuclear matter are 6, 12 and -6 , respectively.

with $M = 939 \text{ MeV}$ the (average) nucleon mass. The next term in this series, $k_f^6/16M^5$, is negligibly small at the densities of interest.

Next, we come to interaction contributions to $S(k_f)$. The closed in-medium diagrams related to one-pion exchange (Fock diagram) and iterated one-pion exchange (Hartree and Fock diagrams) are shown in Fig. 1. Differences in comparison to the calculation of the energy per particle $\bar{E}(k_f)$ in ref.[2] occur only with respect to the factors emerging from the spin-traces over closed nucleon lines and the radii $k_{\uparrow,\downarrow} = k_f(1 \pm \eta)^{1/3}$ of the Fermi spheres to be integrated over. After some analytical calculation we find the following contribution to the spin-asymmetry energy $S(k_f)$ from the 1π -exchange Fock diagram in Fig.1 (including its relativistic $1/M^2$ -correction):

$$S(k_f) = \frac{g_A^2 m_\pi^3}{(4\pi f_\pi)^2} \left\{ \left(\frac{u}{3} + \frac{1}{8u} \right) \ln(1 + 4u^2) - \frac{u}{2} - \frac{u^3}{3} + \frac{m_\pi^2}{M^2} \left[\frac{5u^3}{6} - \frac{u^2}{2} \arctan 2u + \frac{u}{24} (1 - 6u^2) \ln(1 + 4u^2) \right] \right\}. \quad (4)$$

Here, we have introduced the useful abbreviation $u = k_f/m_\pi$ where $m_\pi = 135 \text{ MeV}$ stands for the (neutral) pion mass. As usual $f_\pi = 92.4 \text{ MeV}$ denotes the weak pion decay constant and we choose the value $g_A = 1.3$ of the nucleon axial vector coupling constant in order to have a pion-nucleon coupling constant of $g_{\pi N} = g_A M/f_\pi = 13.2$ [12]. In the second and third diagram in Fig. 1 the 1π -exchange interaction is iterated (once) with itself. These second order diagrams carry the large scale enhancement factor M (the nucleon mass). It stems from an energy denominator that is equal to a difference of small nucleon kinetic energies. With a medium insertion at each of two equally oriented nucleon propagators we obtain from the three-loop Hartree diagram in Fig. 1 the following contribution to the spin-asymmetry energy:

$$S(k_f) = \frac{\pi g_A^4 M m_\pi^4}{6(4\pi f_\pi)^4} \left\{ \left(15u + \frac{7}{2u} \right) \ln(1 + 4u^2) - 14u - 16u^2 \arctan 2u \right\}. \quad (5)$$

The right Fock diagram of iterated 1π -exchange (see Fig. 1) with two medium insertions on non-neighboring nucleon propagators gives rise on the other hand to a contribution to the spin-asymmetry energy of the form:

$$S(k_f) = \frac{\pi g_A^4 M m_\pi^4}{3(4\pi f_\pi)^4} \left\{ \frac{32u^3}{15} + \frac{7u}{5} + \left(\frac{11}{10u} + \frac{2u}{3} \right) \ln(1 + u^2) - \left(\frac{1}{u} + 2u \right) \ln(1 + 4u^2) + \left(\frac{32u^4}{15} - 3 \right) \arctan u + (3 + 4u^2) \left[\arctan 2u + \int_0^u dx \frac{\arctan x - \arctan 2x}{u(1 + 2x^2)} \right] \right\}. \quad (6)$$

This expression does not include the contribution of a linear divergence $\int_0^\infty dl$ of the momentum-space loop-integral. In dimensional regularization such a linear divergence is set to zero, whereas in cut-off regularization it is equal to a momentum space cut-off Λ . The additional term specific for cut-off regularization will be given in eq.(13). An in-medium diagram with three medium insertions represents Pauli-blocking effects in intermediate NN-states induced by the filled Fermi sea of nucleons. The unequal filling of the spin-up and spin-down Fermi seas shows its consequences in the spin-asymmetry energy. After some extensive algebraic manipulations we end up with the following double-integral representation of the contribution to the spin-asymmetry energy $S(k_f)$ from the Hartree diagram in Fig. 1 with three medium insertions:

$$\begin{aligned}
S(k_f) = & \frac{g_A^4 M m_\pi^4}{(4\pi f_\pi)^4 u^3} \int_0^u dx x^2 \int_{-1}^1 dy \left\{ \left[\frac{2uxy(3u^2 - 5x^2y^2)}{(u^2 - x^2y^2)} - (u^2 + 5x^2y^2)H \right] \right. \\
& \times \left[\frac{2s^2 + s^4}{1 + s^2} - 2\ln(1 + s^2) \right] + \frac{4u^2 H s^5(4s' - 3s)}{3(1 + s^2)^2} + [2uxy + (u^2 - x^2y^2)H] \\
& \left. \times [(5 + s^2)(3s^2 - 8ss' + 8s'^2) + 8s(1 + s^2)(s'' - 5s' + 3s)] \frac{s^4}{3(1 + s^2)^3} \right\}, \quad (7)
\end{aligned}$$

where we have introduced several auxiliary functions:

$$H = \ln \frac{u + xy}{u - xy}, \quad s = xy + \sqrt{u^2 - x^2 + x^2y^2}, \quad s' = u \frac{\partial s}{\partial u}, \quad s'' = u^2 \frac{\partial^2 s}{\partial u^2}. \quad (8)$$

Note that eq.(7) stems from a nine-dimensional principal-value integral over the product of three Fermi spheres of varying radii $k_{\uparrow,\downarrow} = k_f(1 \pm \eta)^{1/3}$ which has been differentiated twice with respect to η at $\eta = 0$. Of similar structure is the contribution to $S(k_f)$ from the iterated 1π -exchange Fock diagram with three medium insertions. Because of the two different pion propagators in the Fock diagram one ends up (partially) with a triple-integral representation for its contribution to the spin-asymmetry energy:

$$\begin{aligned}
S(k_f) = & \frac{g_A^4 M m_\pi^4}{24(4\pi f_\pi)^4 u^3} \int_0^u dx \left\{ G(3G_{20} - 2G_{11} + 3G_{02} - 8G_{01} - 3G) \right. \\
& + 3G_{10}^2 - 2G_{01}G_{10} + G_{01}^2 + 4x^2 \int_{-1}^1 dy \int_{-1}^1 dz \frac{yz \theta(y^2 + z^2 - 1)}{|yz| \sqrt{y^2 + z^2 - 1}} \\
& \times \left[\frac{2s^3 t^3 (8s't - 3st - 4s't')}{(1 + s^2)(1 + t^2)} + \frac{s^2[t^2 - \ln(1 + t^2)]}{(1 + s^2)^2} \right] \\
& \left. \times [(3 + s^2)(8ss' - 3s^2 - 8s'^2) + 4s(1 + s^2)(6s' - 3s - 2s'')] \right\}. \quad (9)
\end{aligned}$$

Here, we have split into "factorizable" and "non-factorizable" parts. These two pieces are distinguished by whether the (remaining) nucleon propagator in the three-loop Fock diagram can be canceled or not by terms from the product of πN -interaction vertices. The factorizable terms can be expressed through the auxiliary function:

$$G = u(1 + u^2 + x^2) - \frac{1}{4x} [1 + (u + x)^2][1 + (u - x)^2] \ln \frac{1 + (u + x)^2}{1 + (u - x)^2}, \quad (10)$$

and its partial derivatives for which we have introduced a (short-hand) double-index notation:

$$G_{ij} = x^i u^j \frac{\partial^{i+j} G}{\partial x^i \partial u^j}, \quad 1 \leq i + j \leq 2. \quad (11)$$




Figure 2: The spin-asymmetry energy $S(k_f)$ of nuclear matter versus the nucleon density $\rho = 2k_f^3/3\pi^2$. The full line shows the result of a calculation up to fourth order in small momenta including 1π -exchange and iterated 1π -exchange. The cut-off scale $\Lambda = 0.61$ GeV has been adjusted to the saturation point: $\rho_0 = 0.173 \text{ fm}^{-3}$, $\bar{E}(k_{f0}) = -15.3$ MeV. The isospin-asymmetry energy has the value $A(k_{f0}) = 38.9$ MeV at saturation density. The negative values of $S(k_f)$ indicate the spin-instability of nuclear matter in this approximation.

For the presentation of the nonfactorizable terms one needs also copies of the quantities s and s' defined in eq.(8) which depend (instead of y) on another directional cosine z :

$$t = xz + \sqrt{u^2 - x^2 + x^2 z^2}, \quad t' = u \frac{\partial t}{\partial u}. \quad (12)$$

For the numerical evaluation of the $dydz$ -double integral in eq.(9) it is advantageous to first antisymmetrize the integrand both in y and z and then to substitute $z = \sqrt{y^2 \zeta^2 + 1 - y^2}$. This way the integration region becomes equal to the unit-square $0 < y, \zeta < 1$. In the chiral limit $m_\pi = 0$ the fourth order contributions in eqs.(5-9) sum up to a negative k_f^4 -term of the form: $S(k_f)|_{m_\pi=0} = (g_A k_f / 4\pi f_\pi)^4 (M/135)(16\pi^2 + 273 - 936 \ln 2)$. The formulas for the contributions to the spin-asymmetry energy $S(k_f)$ written down here in eqs.(3-9) have a strong similarity with the analogous contributions to the isospin-asymmetry energy $A(k_f)$ collected in eqs.(20-26) of ref.[2]. This is not surprising in view of the spin-isospin structure $\vec{\sigma} \cdot \vec{\nabla} \tau_a$ of the pion-nucleon coupling. Finally, we give the expression for the linear divergence specific to cut-off regularization:

$$S(k_f) = -\frac{2g_A^4 M \Lambda}{(4\pi f_\pi)^4} k_f^3, \quad (13)$$

to which only the iterated 1π -exchange Fock diagram (with two medium insertions) has contributed. In the case of the Hartree diagram the linear divergence drops out after taking the second derivative with respect to η .

Now we can turn to numerical results. In Fig. 2 we show the spin-asymmetry energy $S(k_f)$ of nuclear matter as a function the nucleon density $\rho = 2k_f^3/3\pi^2$. The full line corresponds to a calculation up to fourth order in small momenta. It includes besides the kinetic energy term eq.(3) the contributions from static 1π -exchange and iterated 1π -exchange. For reasons of consistency we have dropped the small relativistic $1/M^2$ -correction in eq.(4) since it is of fifth order in the small momenta k_f and m_π . The cut-off scale $\Lambda = 611$ MeV has been adjusted to the nuclear matter saturation point $\rho_0 = 0.173 \text{ fm}^{-3}$ and $\bar{E}(k_{f0}) = -15.3$ MeV [7]. The value of the nuclear matter compressibility, $K = k_{f0}^2 \bar{E}''(k_{f0}) = 252$ MeV, as predicted in this framework is consistent with a recent extrapolation from giant monopole resonances of heavy nuclei in ref.[13], which gave $K = (260 \pm 10)$ MeV. One can read off from Fig. 2 a negative value of the spin-asymmetry energy at saturation density: $S(k_{f0}) = S(2m_\pi) = -20.1$ MeV. It reveals the spin-instability of nuclear matter in this restricted framework. The small positive values of $S(k_f)$ at very low densities $\rho \leq 0.03 \text{ fm}^{-3}$ result from the incomplete cancellation between the kinetic energy term eq.(3) proportional to k_f^2 and interaction contributions which grow in magnitude like k_f^3 (or higher powers of k_f). The largest negative contribution to $S(2m_\pi) = -20.1$ MeV comes from the term linear in the cut-off Λ , eq.(13), which amounts to -35.5 MeV at saturation density $k_{f0} = 2m_\pi$. Despite this overwhelming negative contribution to $S(k_f)$ one should not jump to the conclusion that the use of cut-off regularization were the reason for the spin-instability of nuclear matter in this calculation to fourth order. In essence the cut-off Λ is merely parameterizing the strength of a zero-range (S-wave) NN-contact interaction with a fixed ratio of its total-isospin $I = 0$ and $I = 1$ components. The isospin factor $(\vec{\tau}_1 \cdot \vec{\tau}_2)^2 = 3 - 2\vec{\tau}_1 \cdot \vec{\tau}_2$ of the iterated 1π -exchange diagram determines this ratio as 9 : 1. The quality of such a fixed ratio can be tested via the isospin-asymmetry energy $A(k_f)$ (for explicit expressions see section 3 in ref.[2]). The predicted value $A(k_{f0}) = 38.9$ MeV for the isospin-asymmetry energy at saturation density of the present fourth order calculation employing cut-off regularization agrees within 10% with a recent empirical determination, $A(k_{f0}) = (34 \pm 2)$ MeV [13]. The prescribed ratio 9 : 1 between the $I = 0$ and $I = 1$ components of the emerging contact-interaction works therefore rather well. A key observation in this context is that terms linear in the density ρ satisfy the relation: $3S(k_f)_{\text{lin}} + 3A(k_f)_{\text{lin}} + 2\bar{E}(k_f)_{\text{lin}} = 0$. From that one can conclude that even with two free parameters in the contact-interaction (adjusted to the empirical values of $\bar{E}(k_{f0})$ and $A(k_{f0})$) nuclear matter will be spin-unstable in a calculation restricted to fourth order in small momenta. For example, in dimensional regularization where the linear divergence $\int_0^\infty dl 1$ is set to zero one would instead employ a two-parameter momentum-independent NN-contact interaction. The adjustment to the empirical nuclear matter saturation point $\bar{E}(k_{f0} = 2m_\pi) = -15.3$ MeV introduces then the explicit term linear in density $\bar{E}(2m_\pi)_{\text{lin}} = -177.4$ MeV while the constraint from the empirical isospin-asymmetry energy $A(k_{f0}) = 34$ MeV requires in addition $A(2m_\pi)_{\text{lin}} = 148.9$ MeV. These fixed contributions imply via the abovementioned relation a term linear in density for the spin-asymmetry energy of $S(2m_\pi)_{\text{lin}} = -30.6$ MeV. It is almost as negative as the value -35.5 MeV obtained in cutoff regularization with only one adjusted cut-off scale Λ and therefore the spin-instability of nuclear matter would still remain.

The foregoing discussion has shown that it is mandatory to include contributions of fifth order in the small momentum expansion in order to be able to reach the spin-stability of nuclear matter. In the three-loop approximation the terms of fifth order are generated by (irreducible) two-pion-exchange between nucleons. The corresponding one-loop diagrams for elastic NN-scattering are shown in Fig. 3. Since we are counting the delta-nucleon mass splitting $\Delta = 293$ MeV (together with k_f and m_π) as a small momentum scale the diagrams with single and double virtual $\Delta(1232)$ -isobar excitation shown in Fig. 4 belong to the same order. The non-relativistic nucleon and delta propagators are both counted in this scheme as an inverse




Figure 3: One-loop diagrams of irreducible two-pion exchange between nucleons.




Figure 4: One-loop two-pion exchange diagrams with single and double $\Delta(1232)$ -isobar excitation. Diagrams for which the role of both nucleons is interchanged are not shown.

small momentum.

By closing the two open nucleon lines of the one-loop diagrams in Figs.3,4 to either two or one ring one gets (in diagrammatic representation) the Hartree or Fock contribution to the energy density of nuclear matter. The Hartree contribution to the spin-asymmetry energy $S(k_f)$ vanishes identically because the relevant 2π -exchange NN T-matrix in forward direction is spin-independent [4, 14]. The Fock contribution on the other hand is obtained by integrating the spin- and isospin-contracted T-matrix over the product of two Fermi spheres of radii $k_{f,\uparrow,\downarrow} = k_f(1 \pm \eta)^{1/3}$. We separate regularization dependent short-range contributions to the T-matrix (originating from the ultra-violet divergences of the one-loop diagrams in Figs.3,4) from the unique long-range terms with the help of a twice-subtracted dispersion relation. The occurring subtraction constants give rise to a contribution to the spin-asymmetry energy of the form:

$$S(k_f) = (B_3 - 6B_{n,3}) \frac{k_f^3}{3M^2} + S_5 \frac{k_f^5}{M^4}. \quad (14)$$

The two dimensionless parameters $B_3 = -7.99$ and $B_{n,3} = -0.95$ have been adjusted in ref.[3] to few empirical nuclear matter properties (such as the maximal binding energy per particle $-\bar{E}(k_{f0}) = 16$ MeV and the isospin-asymmetry energy $A(k_{f0}) = 34$ MeV with $k_{f0} = 261.6$ MeV). Again, we recognize in the first part of eq.(14) the relation $3S(k_f)_{\text{lin}} + 3A(k_f)_{\text{lin}} + 2\bar{E}(k_f)_{\text{lin}} = 0$ for terms linear in density $\rho = 2k_f^3/3\pi^2$. The other subtraction constant S_5 in front of the k_f^5/M^4 -term is (a priori) not constrained by any property of spin-saturated nuclear matter. The long-range parts of the 2π -exchange (two-body) Fock diagrams can be expressed as a dispersion-integral:

$$S(k_f) = \frac{1}{36\pi^3} \int_{2m_\pi}^{\infty} d\mu \left\{ \text{Im}(V_C + 3W_C) \left[3\mu k_f - \frac{6k_f^3}{\mu} + \frac{16k_f^5}{\mu^3} - \frac{3\mu^3}{4k_f} \ln \left(1 + \frac{4k_f^2}{\mu^2} \right) \right] \right. \\ \left. + \text{Im}(V_T + 3W_T) \left[6\mu^3 k_f + 4\mu k_f^3 - \frac{\mu^3}{2k_f} (8k_f^2 + 3\mu^2) \ln \left(1 + \frac{4k_f^2}{\mu^2} \right) \right] \right\}, \quad (15)$$




Figure 5: Hartree and Fock three-body diagrams related to 2π -exchange with single virtual Δ -isobar excitation. They represent interactions between three nucleons in the Fermi sea. In the case of isospin-symmetric nuclear matter the isospin factors of these diagrams are 8, 0, and 8, in the order shown. The combinatoric factor is 1 for each diagram.

where $\text{Im}V_C$, $\text{Im}W_C$, $\text{Im}V_T$ and $\text{Im}W_T$ are the spectral functions of the isoscalar and isovector central and tensor NN-amplitudes, respectively. Explicit expressions of these imaginary parts for the contributions of the triangle diagram with single Δ -excitation and the box diagrams with single and double Δ -excitation can be easily constructed from the analytical formulas given in section 3 of ref.[4]. The μ - and k_f -dependent weighting functions in eq.(15) take care that at low and moderate densities this spectral-integral is dominated by low invariant $\pi\pi$ -masses $2m_\pi < \mu < 1\text{ GeV}$. The contributions to the spin-asymmetry energy $S(k_f)$ from irreducible 2π -exchange (with only nucleon intermediate states, see Fig. 3) can also be cast into the form eq.(15). The corresponding non-vanishing spectral functions read [14]:

$$\text{Im}W_C(i\mu) = \frac{\sqrt{\mu^2 - 4m_\pi^2}}{3\pi\mu(4f_\pi)^4} \left[4m_\pi^2(1 + 4g_A^2 - 5g_A^4) + \mu^2(23g_A^4 - 10g_A^2 - 1) + \frac{48g_A^4 m_\pi^4}{\mu^2 - 4m_\pi^2} \right], \quad (16)$$

$$\text{Im}V_T(i\mu) = -\frac{6g_A^4 \sqrt{\mu^2 - 4m_\pi^2}}{\pi\mu(4f_\pi)^4}. \quad (17)$$

Next, we come to the additional 2π -exchange three-body terms which arise from Pauli blocking of intermediate nucleon states (i.e. from the $-2\pi\theta(k_{\uparrow,\downarrow} - |\vec{p}|)$ terms in the in-medium nucleon propagators [2]). The corresponding closed Hartree and Fock diagrams with single virtual Δ -excitation are shown in Fig. 5. For isospin-symmetric nuclear matter their isospin factors are 8, 0, and 8, in the order shown. The contribution of the left three-body Hartree diagram to the spin-asymmetry energy $S(k_f)$ has the following analytical form:

$$S(k_f) = \frac{g_A^4 m_\pi^6 u^2}{9\Delta(2\pi f_\pi)^4} \left[\left(\frac{9}{4} + 4u^2 \right) \ln(1 + 4u^2) - 2u^4(1 + 3\zeta) - 8u^2 - \frac{u^2}{1 + 4u^2} \right]. \quad (18)$$

The delta propagator shows up in this expression merely via the (reciprocal) mass-splitting $\Delta = 293\text{ MeV}$. Furthermore, we have already inserted in eq.(18) the empirically well-satisfied relation $g_{\pi N\Delta} = 3g_{\pi N}/\sqrt{2}$ for the $\pi N\Delta$ -coupling constant. The parameter $\zeta = -3/4$ has been introduced in section 2 of ref.[3] in order to reduce a too strongly repulsive ρ^2 -term in the energy particle $\bar{E}(k_f)$. It controls the strength of a three-nucleon contact interaction $\sim (\zeta g_A^4/\Delta f_\pi^4)(NN)^3$ which has the interesting property that it contributes equally but with opposite sign to the energy per particle $\bar{E}(k_f)$ and the spin-asymmetry energy $S(k_f)$. The contribution of the right three-body Fock diagram in Fig. 5 to the spin-asymmetry energy $S(k_f)$ can be represented as:

$$S(k_f) = \frac{g_A^4 m_\pi^6}{36\Delta(4\pi f_\pi)^4 u^3} \int_0^u dx \left\{ 4G_{S01}G_{S10} - 2G_{S01}^2 - 6G_{S10}^2 \right\}$$




Figure 6: The spin-asymmetry energy $S(k_f)$ of nuclear matter versus the nucleon density $\rho = 2k_f^3/3\pi^2$. In comparison to Fig. 2 the effects from 2π -exchange with single and double virtual Δ -isobar excitation are now included. The full, dashed and dashed-dotted curves correspond to the choices $S_5 = 0, -7.45$ and -30 of the short-range parameter S_5 . The positive values of $S(k_f)$ ensure the spin-stability of nuclear matter.

$$+2G_S(3G_S + 8G_{S01} - 3G_{S02} + 2G_{S11} - 3G_{S20}) + 2G_{T01}G_{T10} - 7G_{T01}^2 - 3G_{T10}^2 + G_T(3G_T + 8G_{T01} - 3G_{T02} + 2G_{T11} - 3G_{T20})\}, \quad (19)$$

with the two auxiliary functions:

$$G_S = \frac{4ux}{3}(2u^2 - 3) + 4x \left[\arctan(u+x) + \arctan(u-x) \right] + (x^2 - u^2 - 1) \ln \frac{1 + (u+x)^2}{1 + (u-x)^2}, \quad (20)$$

$$G_T = \frac{ux}{6}(8u^2 + 3x^2) - \frac{u}{2x}(1 + u^2)^2 + \frac{1}{8} \left[\frac{(1 + u^2)^3}{x^2} - x^4 + (1 - 3u^2)(1 + u^2 - x^2) \right] \ln \frac{1 + (u+x)^2}{1 + (u-x)^2}. \quad (21)$$

The double-indices on G_S and G_T have the same meaning as explained in eq.(11) for the function G . It is remarkable that the three-body Hartree and Fock diagrams with virtual Δ -excitation (shown in Fig. 5) lead both to identical contributions to the spin-asymmetry energy $S(k_f)$ and the isospin-asymmetry energy $A(k_f)$ (compare eqs.(18,19) here with eqs.(48,49) in ref.[3]). This feature originates from the specific spin-isospin structure of the (non-relativistic) $\pi N \Delta$ -transition vertex [4].

In Fig. 6 we show again the spin-asymmetry energy $S(k_f)$ of nuclear matter as a function of the nucleon density $\rho = 2k_f^3/3\pi^2$. The full line includes all the contributions from chiral 1π - and

2π -exchange written down in eqs.(3-9,14-19). The (yet undetermined) short-range parameter S_5 has been set to zero, $S_5 = 0$. We note as an aside that the term linear in the density and the cut-off Λ eq.(13) is of course now not counted extra since the parameter $B_3 - 6B_{n,3} = -2.29$ [3] collects all such possible terms. One observes in Fig.6 a positive spin-asymmetry energy $S(k_f)$ which rises monotonically with the density ρ . The inclusion of the chiral $\pi N\Delta$ -dynamics has made nuclear matter spin-stable. This result is of utmost importance since it assures that the improved nuclear matter properties reported in ref.[3] are based on a spin-stable ground state. It is also interesting to look at numerical values of $S(k_f)$ and their decomposition. At a Fermi momentum of $k_f = 2m_\pi$ (corresponding to $\rho = 0.173 \text{ fm}^{-3}$) the spin-asymmetry energy is now $S(2m_\pi) = 67.3 \text{ MeV}$ (setting $S_5 = 0$). The most significant changes in comparison to the previous fourth order calculation come from the term linear in density which gets reduced (by a factor of 2.1) to -17.0 MeV and the two-body Fock and three-body Hartree contributions eqs.(15,18) which amount together to $54.6 \text{ MeV} + 13.7 \text{ MeV} = 68.3 \text{ MeV}$. We note also that about one quarter thereof (namely 16.2 MeV) stem from the three-body contact interaction proportional to $\zeta = -3/4$.

The remaining open question concerns the size of the short-distance parameter S_5 in eq.(14). Large negative values could again endanger the spin-stability of nuclear matter. In order to get an estimate of S_5 we bring into play the complete set of four-nucleon contact-couplings written down in eqs.(3,4) of ref.[15]. This set represents the most general short-range NN-interaction quadratic in momenta and it involves seven low-energy constants $C_1, \dots, C_7$. After computing the spin-asymmetry energy $S(k_f)$ from the corresponding contact-potential in Hartree-Fock approximation we find:

$$\begin{aligned} S_5 &= \frac{M^4}{18\pi^2} (C_2 - 4C_1 + 3C_4 + C_7) \\ &= \frac{M^4}{96\pi^3} \left[2C(^3S_1) - 2C(^1S_0) - C(^1P_1) + C(^3P_0) + 3C(^3P_1) + 5C(^3P_2) \right]. \end{aligned} \quad (22)$$

In the second line of eq.(22) we have reexpressed the relevant linear combination of $C_{1,2,4,7}$ through the so-called spectroscopic low-energy constants which characterize the short-range part of the NN-potential in the spin-singlet and spin-triplet S - and P -wave states. In that representation we obtain from the entries of table IV in ref.[15] for the three high-precision NN-potentials CD-Bonn [16], Nijm-II [17] and AV-18 [18] the numbers $S_5 = -7.10, -7.98$ and -7.27 . The dashed line in Fig.6 shows the spin-asymmetry energy $S(k_f)$ which results from taking their average value $S_5 = -7.45$. The effective reduction of the spin-asymmetry energy by about 20% is harmless and the spin-stability of nuclear matter is equally well guaranteed. The dashed-dotted curve in Fig.6 corresponds to the extreme choice $S_5 = -30$. One can see that even with such a large negative S_5 -value, which exceeds the estimate from realistic NN-potentials by a factor of 4, the spin-stability of nuclear matter still remains preserved. We can therefore conclude that with the inclusion of the chiral $\pi N\Delta$ -dynamics spin-stability becomes a robust property of nuclear matter. This is an important finding and it goes conform with the improved isospin properties observed in ref.[3].

In summary we have investigated in this work the spin-stability of nuclear matter in the framework of chiral perturbation theory. For that purpose we have calculated the density-dependent spin-asymmetry energy $S(k_f)$ of isospin-symmetric nuclear matter to three-loop order. The interaction contributions to $S(k_f)$ originate from one-pion exchange, iterated one-pion exchange, and (irreducible) two-pion exchange with no, single, and double virtual Δ -isobar excitation. We have found that the truncation to 1π -exchange and iterated 1π -exchange terms (which leads already to a good nuclear matter equation of state) is spin-unstable, since

$S(k_{f0}) < 0$. This statement holds independently of the regularization scheme if the contact terms (generating contributions linear in density) are consistent with the empirical nuclear matter bulk properties: $\bar{E}(k_{f0}) \simeq -16$ MeV and $A(k_{f0}) \simeq 34$ MeV. The inclusion of the chiral $\pi N\Delta$ -dynamics on the other hand guarantees the spin-stability of nuclear matter. The corresponding spin-asymmetry energy $S(k_f)$ stays positive within a wide range of an undetermined short-range parameter S_5 , which we have also estimated from realistic NN-potentials. Our results reemphasize the important role played by two-pion exchange with virtual Δ -isobar excitation for the nuclear matter many-body problem.

References

- [1] M. Lutz, B. Friman, and Ch. Appel, *Phys. Lett.* **B474**, 7 (2000).
- [2] N. Kaiser, S. Fritsch and W. Weise, *Nucl. Phys.* **A697**, 255 (2002); and references therein.
- [3] S. Fritsch, N. Kaiser, and W. Weise, "Chiral approach to nuclear matter: Role of two-pion exchange with virtual delta-isobar excitation", nucl-th/0406038; submitted to *Nucl. Phys.* **A**.
- [4] N. Kaiser, S. Gerstendörfer and W. Weise, *Nucl. Phys.* **A637**, 395 (1998).
- [5] N. Kaiser, S. Fritsch and W. Weise, *Nucl. Phys.* **A700**, 343 (2002).
- [6] S. Fritsch, N. Kaiser and W. Weise, *Phys. Lett.* **B545**, 73 (2002).
- [7] S. Fritsch and N. Kaiser, *Eur. Phys. J.* **A17**, 11 (2003).
- [8] B.D. Chang, *Phys. Lett.* **B56**, 205 (1975).
- [9] S.O. Bäckman, A.D. Jackson and J. Speth, *Phys. Lett.* **B56**, 209 (1975).
- [10] S. Krewald, V. Klemt, J. Speth and A. Faessler, *Nucl. Phys.* **A281**, 166 (1977).
- [11] F.L. Braghin, *Int. J. Mod. Phys.* **E12**, 755 (2003); and references therein.
- [12] M.M. Pavan et al., *Phys. Scr.* **T87**, 65 (2000).
- [13] D. Vretenar, T. Niksic and P. Ring, *Phys. Rev.* **C68**, 024310 (2003).
- [14] N. Kaiser, R. Brockmann and W. Weise, *Nucl. Phys.* **A625**, 758 (1997).
- [15] E. Epelbaum, Ulf-G. Meißner, W. Glöckle, and C. Elster, *Phys. Rev.* **C65**, 044001 (2002); and references therein.
- [16] R. Machleidt, *Phys. Rev.* **C63**, 024001 (2001); and references therein.
- [17] V.G.J. Stoks, R.A.M. Klomp, C.P.F. Terheggen and J.J. de Swart, *Phys. Rev.* **C49**, 2950 (1994).
- [18] R.B. Wiringa, V.G.J. Stoks and R. Schiavilla, *Phys. Rev.* **C51**, 38 (1995).